

## **Anti C-Peptide II (Rat) Serum**

Cat. No. Y220

Lot No. 90170927

**Description:** This antiserum was raised in a rabbit by immunization with a carrier free synthetic C-peptide II (rat) peptide. The product vial contains 50  $\mu$ L of the titled antiserum, which was obtained by lyophilizing its 0.001M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with C-peptide (rat).

**Immunogen:** Synthetic C-peptide II (rat), carrier free

**Host:** Rabbit

**Amino Acid Sequence of C-peptide II (rat)<sup>1,2)</sup>**

EVEDPQVAQL ELGGGPGAGD LQTLALEVAR Q

**Product Form:** Lyophilized unpurified serum

**Size:** 50  $\mu$ L

**Reconstitution:** Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN<sub>3</sub> 0.1%).

**Storage:** The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

**Suggested Working Dilution Range:** 1:1,250 (final dilution 1:10,000) for radioimmunoassay; 1:1,000-10,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

**Specificity (based on RIA):** C-peptide I (rat) 100%, C-peptide II (rat) 100%, C-peptide I (7-31) (rat) 100%, C-peptide II (7-31) (rat) 100%, C-peptide I (15-31) (rat) 100%, C-peptide II (15-31) (rat) 100%, C-peptide I (20-31) (rat) 80 %, C-peptide I (1-23) (rat) <0.001%, C-peptide II (1-23) (rat) <0.001%, C-peptide (human) 0.01%<sup>3)</sup>

**Positive Control (immunohistochemistry):** Rat pancreas.


**Species Tested:** Rat

### **REFERENCES:**

- 1) J. Markussen and F. Sundby, Rat-proinsulin C-peptides. Amino-acid sequences, *Europ J Biochem*, 25: 153, 1972
- 2) H.S. Tager and D.F. Steiner. Primary Structures of the proinsulin connecting peptides of the rat and the horse, *J Biol Chem*, 247:7936-7940, 1972
- 3) C. Yanaihara, J. Ozaki et al., Immunological studies on synthetic rat and guinea pig C-peptides. In: *Proinsulin, Insulin, C-peptide* (S Baba et al Ed) Excerpta Medica, Amsterdam-Oxford, pp87-93, 1979

**FOR RESEARCH LABORATORY USE ONLY**

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

  
Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan  
TEL: +81-544-22-2771 FAX: +81-544-22-2770

For more information, please visit our internet website: <http://www.yanaihara.co.jp> or contact us to: [ask@yanaihara.co.jp](mailto:ask@yanaihara.co.jp)