

Anti C-Peptide I (Mouse) Serum

Cat. No. Y222

Description: This antiserum was raised in a rabbit by immunization with a keyhole limpet hemocyanin (KLH) conjugate of synthetic C-peptide I (mouse) peptide. The product vial contains 50 μ L of the titled antiserum, which was obtained by lyophilizing its 0.001M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with C-peptide I (mouse).

Immunogen: Synthetic C-peptide I (mouse)-KLH conjugate

Host: Rabbit

Amino Acid Sequence of C-peptide I (mouse)¹⁾

EVEDPQVEQL ELGGSPGDLQ TLALEVARQ

Product Form: Lyophilized unpurified serum

Size: 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing- thawing should be avoided.

Suggested Working Dilution Range: 1:1,000-5,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on non-competitive EIA): C-peptide I (mouse) 100%, C-peptide II (mouse) 15.2%, C-peptide I (rat) 20.9%, C-peptide II (rat) 85%, C-peptide (human) < 0.01%, C-peptide (dog) < 0.01%, C-peptide (porcine) < 0.1%, glucagon < 0.01%,

Positive Control (immunohistochemistry): Mouse pancreas.

Species Tested: Mouse

REFERENCES:

1) B.M. Wentworth, I.M. Schaefer et al., Characterization of the two nonallelic genes encoding mouse preproinsulin. *Journal of Molecular Evolution*, 23: 305-312, 1986

FOR RESEARCH LABORATORY USE ONLY
DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM



For more information, please visit our internet website: <http://www.yanaihara.co.jp> or contact us to ask@yanaihara.co.jp

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan
FAX: +81-544-22-2770 TEL: +81-544-22-2771